



The Relationship between Inotrope Use and Intraventricular Hemorrhage in Preterm Neonates

Sara Mokhtar Abdelaal Hedia ^{a*}, Manal Ezzat Badawy ^b,
Hamed Mohamed Elsharkawy ^a
and Sara Mohamed Elashry ^a

^a Pediatric Department, Faculty of Medicine, Tanta University, Tanta, Egypt.

^b Radiology Department, Faculty of Medicine, Tanta University, Tanta, Egypt.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajpr/2024/v14i12413>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/128019>

Review Article

Received: 14/10/2024

Accepted: 16/12/2024

Published: 27/12/2024

ABSTRACT

In preterm newborns, intraventricular haemorrhage (IVH) may be preceded by erroneous hemodynamic alterations in cerebral circulation. Neonatal hypotension requiring use of inotropes was found to be the significantly risk factors in IVH. Reduced use of inotropic medications (such dopamine or epinephrine), which have been linked to the development of IVH, and a reduced incidence of hypotension, may be linked to the increased intravascular volume.

Keywords: Inotrope; intraventricular hemorrhage; preterm neonates.

*Corresponding author: Email: sara.elashry@med.tanta.edu.eg;

Cite as: Hedia, Sara Mokhtar Abdelaal, Manal Ezzat Badawy, Hamed Mohamed Elsharkawy, and Sara Mohamed Elashry. 2024. "The Relationship Between Inotrope Use and Intraventricular Hemorrhage in Preterm Neonates". *Asian Journal of Pediatric Research* 14 (12):84-88. <https://doi.org/10.9734/ajpr/2024/v14i12413>.

1. INTRODUCTION

IVH is the most prevalent and serious neurological disorder linked to prematurity, impacting 20–40% of all infants born globally who weigh less than 1500 g. It is believed that neonatal IVH occurs due to the interplay between delicate germinal matrix vessels and the swift hemodynamic fluctuations that premature infants are susceptible to. Newborns may undergo ischemia as a result of decreased heart rate, which is subsequently followed by reperfusion as ventricular function starts to recover during the initial days of life (Deger et al., 2021).

These abrupt instances of ischemia-reperfusion can place stress on the vessels of the germinal matrix, resulting in IVH (Ballabh & de Vries, 2021; Lai et al., 2022). It is believed that this phenomenon happens in premature infants because of an immature germinal matrix, developing brain autoregulatory mechanisms, and heightened hemodynamic variations linked to underdeveloped cardiopulmonary systems (Deger et al., 2021).

Factors such as the underdevelopment of autoregulatory mechanisms predispose premature infants, leading to decreased cerebral blood flow as a result of variations in MAP and carbon dioxide partial pressure (Stathopoulou et al., 2023).

Inotropes are drugs that alter the contractile force of the heart muscle. Contractility is increased by positive inotropes and decreased by negative inotropes. These chemicals affect smooth muscle in both excitatory and inhibitory ways of the heart and blood vessels, as well as important effects on metabolism, the presynaptic autonomic nervous system and the central nervous system (Grandmougin et al., 202; Scheeren et al., 2021; Darweesh et al., 2022; Elsheikh et al., 2024). They are frequently used to increase vascular tone (RVS) or cardiac output (CO) in order to enhance haemodynamics to provide oxygen to sustain important organs while the fundamental illness is being addressed (Motwani & Saunders, 2021).

2. INTRAVENTRICULAR HEMORRHAGE

Premature babies are susceptible to periventricular-intraventricular haemorrhage (PIVH). When the arteries in the periventricular

region's germinal matrix burst, haemorrhage occurs, which may proceed into the ventricles as intraventricular haemorrhage (IVH). In extreme cases, the bleeding will expand inhabit a sizable section of the ventricle and extend into the intraparenchymal region. Due to the fact that the germinal matrix begins to involute during 33 weeks gestation, infants born before this time are the most vulnerable (Gotardo et al., 2019).

The global prevalence of peripheral viral hepatitis ranges from 3.70% to 44.68%. Studies have revealed a 36.2% overall incidence, with 7% of cases exhibiting severe grades (III, IV). The overall prevalence of class I, II, III, and IV PIVH in preterm infants is 17.0%, 12.1%, 3.3%, and 3.8%, respectively, with 1% of cases (van Bel & Mintzer, 2018).

Risk factors: Germinal matrix-IVH is mostly caused by extremely low gestational age. Pregnant women who give birth before 32 weeks are considered high-risk. Asphyxia, respiratory distress, pneumothorax, pulmonary haemorrhage, choroid amnionitis, mechanical ventilation, sepsis, patent ductus arteriosus, and variations in cerebral blood flow (Du et al., 2022).

It was shown that elevations in arterial blood pressure and the incidence of IVH were strongly correlated with changes in cerebral blood flow. Reduced haematocrit levels, hypercarbia, fast volume replacements, hypoglycemia, and disruption of cerebral autoregulation are the primary reasons for the increase in cerebral blood flow. In premature newborns with abnormally low birth weights, hypernatraemia was found to be an independent risk factor for IVH (Dalton et al.).

Cerebral blood flow fluctuation: Cerebrovascular autoregulation (CAR) is crucial for ensuring consistent and sufficient cerebral blood flow during hypotension, hypoperfusion, or hypertension (Klein et al., 2019).

Hypotension and hypertension: Irregular hemodynamic fluctuations in cerebral circulation may occur before IVH in preterm infants (Tsao, 2023).

Preterm infants with hypotension underwent notably extended durations of cerebral hypoxia and had impaired CAR, with these traits linked to early IVH or mortality (Thewissen et al., 2021).

The incidence of IVH was shown to be higher in critically sick preterm newborns with pressure-passive circulation than in neonates with good autoregulation. This implies that preterm newborns are more negatively impacted by high blood pressure. A considerably higher frequency of severe IVH was linked to extreme mean arterial blood pressure (MABP) values (i.e., <23 and >46 mmHg) in preterm infants born at a gestational age of less than 30 weeks (Vesoulis et al., 2020).

Inotropes: An inotrope, or inotropic agent, refers to a drug or substance that modifies the strength or energy of muscle contractions. Negatively inotropic agents reduce the strength of muscle contractions. Agents that are positively inotropic enhance the force of muscular contractions (DesJardin & Teerlink, 2021).

Mechanism of action: Drugs known as vasopressors and inotropes cause vasoconstriction or enhance cardiac contractility in shock patients or any other condition that results in abnormally low blood pressure. One of the main features of shock is a decreased blood flow to essential organs, which can result in multiple organ failure and eventually death (Polyzogopoulou et al., 2020).

Vasopressors increase contractility and heart rate, which in turn causes peripheral vasoconstriction, which raises systemic vascular resistance (SVR) and CO. Raising the SVR leads to increased organ perfusion and mean arterial pressure (MAP) (Manaker & Parsons, 2020). By enhancing cardiac contractility, inotropes raise cardiac output (CO), supporting body perfusion and MAP.

The relationship between the two is expressed by the equation $MAP = CO \times SVR$ (Manaker & Parsons, 2020).

- Inotrope associated with intraventricular hemorrhage

Prophylactic indomethacin usage (Foglia et al., 2018), limiting the number of intubation attempts for extremely preterm newborns (St Peter et al., 2017), and reducing the use of inotropes are some of the strategies used to lower the incidence of severe IVH (Abdul et al., 2020).

The elevated intravascular volume might correlate with a lower occurrence reduction in the use of inotropic medications (such as

dopamine or epinephrine) and hypotension, both of which have been linked to the development of IVH (Alotaibi et al., 2020).

Early inotrope use was independently linked to mortality and/or severe brain injury in preterm children delivered before 29 weeks of gestation, and a greater prevalence of both severe and any IVH in infants born between 24 and 27 weeks was associated with the requirement for inotropic treatment within 72 hours of birth.

3. CONCLUSION

Various investigations have shown the independent effect of vasopressors on interrupting cerebral autoregulation, inotropes may be a useful predictor of how serious a patient's condition is (Bouyssi et al., 2018). After adjusting for confounding variables, recent research shown that using inotropes was linked to a higher risk of severe IVH rather than hypotension in and of itself (Alotaibi et al., 2020).

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Abdul Aziz, A. N., Thomas, S., Murthy, P., Rabi, Y., Soraisham, A., Stritzke, A., et al. (2020). Early inotropes use is associated with higher risk of death and/or severe brain injury in extremely premature infants. *J Matern Fetal Neonatal Med*, 33, 2751-2758.
- Alotaibi, W. S. M., Alsaif, N. S., Ahmed, I. A., Mahmoud, A. F., Ali, K., Hammad, A., et al. (2020). Reduction of severe intraventricular hemorrhage, a tertiary single-center experience: Incidence trends, associated risk factors, and hospital policy. *Childs Nerv Syst*, 36, 2971-2979.

- Ballabh, P., & de Vries, L. S. (2021). White matter injury in infants with intraventricular haemorrhage: Mechanisms and therapies. *Nature Reviews Neurology*, 17(4), 199-214.
- Bouyssi-Kobar, M., Murnick, J., Brossard-Racine, M., Chang, T., Mahdi, E., Jacobs, M., et al. (2018). Altered cerebral perfusion in infants born preterm compared with infants born full term. *J Pediatr*, 193, 54-61.
- Darweesh, H. F., Nassar, M. A. E., El Tomy, M. A., & El Mashad, A. R. M. (2022). Pneumothorax in preterm neonates and its association with intraventricular hemorrhage. *Journal of Advances in Medicine and Medical Research*, 34(21), 460-468. <https://doi.org/10.9734/jammr/2022/v34i2131564>
- Deger, J., Goethe, E. A., LoPresti, M. A., & Lam, S. (2021). Intraventricular hemorrhage in premature infants: A historical review. *World Neurosurg*.
- DesJardin, J. T., & Teerlink, J. R. (2021). Inotropic therapies in heart failure and cardiogenic shock: An educational review. *Eur Heart J Acute Cardiovasc Care*, 10, 676-686.
- Du, X., Lei, G., Zhang, J., Liao, C., Chen, X., & Wang, J. (2022). Risk factors of periventricular-intraventricular hemorrhage in premature infants. *Chin Pediatr Emerg Med*, 15, 665-670.
- Elsheikh, R. M., Elmashad, A. M., Aboelenin, A. M., & Elrifay, S. M. (2024). Relative adrenal insufficiency in full-term and preterm neonates with neonatal sepsis. *Asian Journal of Pediatric Research*, 14(1), 1-13. <https://doi.org/10.9734/ajpr/2024/v14i1314>
- Foglia, E. E., Roberts, R. S., Stoller, J. Z., Davis, P. G., Haslam, R., & Schmidt, B. (2018). Effect of prophylactic indomethacin in extremely low birth weight infants based on the predicted risk of severe intraventricular hemorrhage. *Neonatology*, 113, 183-186.
- Gotardo, J. W., Volkmer, N. d. F. V., Stangler, G. P., Dornelles, A. D., Bohrer, B. B. d. A., & Carvalho, C. G. (2019). Impact of periventricular haemorrhage and periventricular leukomalacia in the neurodevelopment of preterms: A systematic review and meta-analysis. *PLoS One*, 14, 427.
- Grandmougin, D., Chalon, A., Falanga, A., Marie, V., Groubatch-Joineau, F., Mourer, B., Marie, P.-Y., Lacolley, P., & Tran, N. (2022). Evidences for inotropic effects induced by simultaneous antegrade and retrograde myocardial perfusion in a swine model: Potential applications for ECMO. *Current Practice in Medical Science*, 5, 188-207. <https://doi.org/10.9734/bpi/cpms/v5/16704D>
- Klein, M. O., Battagello, D. S., Cardoso, A. R., Hauser, D. N., Bittencourt, J. C., & Correa, R. G. (2019). Dopamine: Functions, signaling, and association with neurological diseases. *Cell Mol Neurobiol*, 39, 31-59.
- Lai, G. Y., Shlobin, N., Garcia, R. M., Wescott, A., Kulkarni, A. V., Drake, J., ... & Lam, S. K. (2022). Global incidence proportion of intraventricular haemorrhage of prematurity: A meta-analysis of studies published 2010–2020. *Archives of Disease in Childhood-Fetal and Neonatal Edition*, 107(5), 513-519.
- Manaker, S., Parsons, P., & Finlay, G. (2020). Use of vasopressors and inotropes. *Literature Review Current Through*.
- Manaker, S., Parsons, P., & Finlay, G. (2020). Use of vasopressors and inotropes. *Literature Review Current Through*.
- Motwani, S. K., & Saunders, H. (2021). Inotropes. *Anaesth Intensive Care*, 22, 243-248.
- Polyzogopoulou, E., Arfaras-Melainis, A., Bistola, V., & Parissis, J. (2020). Inotropic agents in cardiogenic shock. *Curr Opin Crit Care*, 26, 403-410.
- Scheeren, T. W., Bakker, J., Kaufmann, T., Annane, D., Asfar, P., Boerma, E. C., ... & Teboul, J. L. (2021). Current use of inotropes in circulatory shock. *Annals of Intensive Care*, 11, 1-13.
- St Peter, D., Gandy, C., & Hoffman, S. B. (2017). Hypotension and adverse outcomes in prematurity: Comparing definitions. *Neonatology*, 111, 228-233.
- Stathopoulou, T., Agakidou, E., Paschaloudis, C., Kontou, A., Chatzioannidis, I., & Sarafidis, K. (2023). Strong association between inotrope administration and intraventricular hemorrhage, gestational age, and the use of fentanyl in very low gestational age infants: A retrospective study. *Children*.
- Thewissen, L., Naulaers, G., Hendriks, D., Caicedo, A., Barrington, K., Boylan, G., et al. (2021). Cerebral oxygen saturation and autoregulation during hypotension in extremely preterm infants. *Pediatr Res*, 90, 373-380.

- Tsao, P. C. (2023). Pathogenesis and prevention of intraventricular hemorrhage in preterm infants. *J Korean Neurosurg Soc*, 66, 228-238.
- van Bel, F., & Mintzer, J. P. (2018). Monitoring cerebral oxygenation of the immature brain: A neuroprotective strategy? *Pediatr Res*, 84, 159-164.
- Vesoulis, Z. A., Flower, A. A., Zanelli, S., Rambhia, A., Abubakar, M., Whitehead, H. V., et al. (2020). Blood pressure extremes and severe IVH in preterm infants. *Pediatr Res*, 87, 69-73.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/128019>