

Journal of Advances in Microbiology

Volume 24, Issue 12, Page 172-179, 2024; Article no.JAMB.128149 ISSN: 2456-7116

Efficacy of Essential Oils against Purple Blotch of Garlic (Allium sativum L.) Caused by Alternaria porri (Ellis) Cif

Sujeet Pandey ^{a++*}, Shashi Tiwari ^{a#} and Yashika Srivastava ^{a++}

^a Department of Plant Pathology, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj-211007, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: https://doi.org/10.9734/jamb/2024/v24i12881

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/128149

Original Research Article

Received: 12/10/2024 Accepted: 14/12/2024 Published: 17/12/2024

ABSTRACT

A research trial was carried out during *Rabi* season, 2023-24 at the Central Research Field, Department of Plant Pathology, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology And Sciences, Prayagraj, Uttar Pradesh. The experiment was conducted to evaluate the effect of different essential oils on purple blotch of garlic caused by *Alternaria porri* on

++M. Sc. Scholar;

Cite as: Pandey, Sujeet, Shashi Tiwari, and Yashika Srivastava. 2024. "Efficacy of Essential Oils Against Purple Blotch of Garlic (Allium Sativum L.) Caused by Alternaria Porri (Ellis) Cif". Journal of Advances in Microbiology 24 (12):172-79. https://doi.org/10.9734/jamb/2024/v24i12881.

[#]Assistant Professor; *Corresponding author: E-mail: sujeet.pandey.2503@gmail.com;

disease intensity and yield. The trial was conducted using seven treatments *viz*. T₁- neem oil (5%), T₂- eucalyptus oil (5%), T₃- clove oil (5%), T₄- castor oil (5%), T₅- mustard oil (5%), T₆- mancozeb 75 WP (0.2%) and T₀- control with each treatment replicating three times to manage the disease and also to evaluate the effect of treatments on the disease intensity and yield. Results showed that all treatments significantly increased yield of garlic and reduced the intensity of disease. From the study of results, it was concluded that among all the treatments, the most effective in managing the disease was T₆- Mancozeb (27.85%) followed by T₁- Neem oil (34.76%), T₃- Clove oil (37.66%), T₂- Eucalyptus oil (37.83%), T₄- Castor oil (39.49%) and T₅- Mustard oil (40.25%) as compared to T₀- Control (49.24%) and for influencing the yield (t/ha.) was T₆- Mancozeb (5.41) followed by T₁- Neem oil (4.05), T₃- Clove oil (3.83), T₂- Eucalyptus oil (3.73), T₄- Castor oil (3.73), T₄- Castor oil (3.56) and T₅- Mustard oil (3.51) as compared to T₀- Control (2.61).

Keywords: Alternaria porri; essential oils; purple blotch; yield.

1. INTRODUCTION

Garlic (Allium sativum L.) (2n=16) is the second most important allium crop after onion grown throughout the plains of India, of family Alliaceae amongst the spices and condiments. It is commonly termed as "Lasan" (Singh et al., 2021). Garlic has higher nutritive value as compared to other bulbous crops. It is a rich source of carbohydrates (29%), proteins (7%), fibres (0.8%), minerals (0.3%), fat (0.2%), essential oils (0.1-0.4 %) and also contains vitamin C and sulphur (Memane et al., 2008). In addition to this, garlic has a wide spectrum of actions. It has antibacterial, antiviral, antifungal and antiprotozoal properties. It is also beneficial to the cardiovascular and immune systems and has antioxidant and anticancer properties (Harris et al., 2001).

Garlic comes from Central Asia, but it's grown all over the world, including in Brazil, Mexico, Spain, India, Egypt, Bulgaria, and Hungary. One of the major bulb crops cultivated and utilized as a spice or condiment all over India is garlic. It foreign contributes significantly to India's exchange earnings. The world's largest producer of garlic is the Chinese mainland. India is the second-largest producer of garlic in the world, with a yield of roughly 81378 kg/ha from 392000 hectares of land, yielding approximately 3190000 tonnes of garlic (FAO, 2021). India's top states for garlic production are Gujarat (3.29%), Uttar Pradesh (6.57%), Madhya Pradesh (62.85%), Rajasthan (16.81%), and Punjab (2.66%) (NHB, 2021-22) (Tripathi and Lawande, 2006).

The major diseases of this crop are purple blotch, botrytis rot, botrytis leaf blight, cercospora leaf spot, downy mildew, fusarium basal rot, damping-off, white rot, stem and bulb nematode, mosaic virus etc. *Alternaria* sp. decay is one of the major factors responsible for economic losses in garlic (Prajapati *et al.*, 2020).

The name purple blotch for this disease was proposed by Nolla (1927). The pathogen of Allium was first designated as Macrosporium porri (Cooke and Ellis, 1879) and Alternaria allii (Nolla, 1927). Later both species were called Alternaria porri (Ciferri, 1930). According to Simmons (2007), the conidia of Alternaria allii were different than those of Alternaria porri based on their multiple branches and beaks. Purple blotch caused by Alternaria porri, is the most destructive disease of Allium sp. (onions, garlic, shallots, leeks, scallions and chives). The pathogenicity of Alternaria sp. is due to production of host specific or nonspecific toxins that may induce disease. These toxins are mainly secondary metabolites that destroy susceptible cultivars by leaf necrosis. The disease usually affects the leaves and bulbs of a plant, and reducing their yield up to 97% (Kareem et al., 2012).

2. MATERIALS AND METHODS

The field investigations were carried out at the Central Research Field of the Department of Plant Pathology, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.) during *Rabi* 2023-24. The seeds were sown in December, 2023 maintaining a distance of 15 cm row to row and 10 cm plant to plant.

2.1 Preparation of Potato Dextrose Agar (PDA) Medium

Potato dextrose agar medium was used for isolation of the pathogen (*Alternaria porri*) from the diseased plant parts and maintenance of cultures. The composition of PDA is as follows:

Peeled potato: 200 g Dextrose: 20 g Agar: 20 g Distilled water: 1000 ml pH: 5.6 - 6.5

200 grams of peeled and cut potatoes were boiled in 1000 ml of distilled water. The potato extract was filtered through muslin cloth and poured in a beaker. 20 grams each of dextrose and agar was dissolved in it, pH was maintained at 5.6- 6.5. The final volume was made up to 1000 ml by adding more of distilled water. In each conical flask 1/3rd of this solution was then dispensed and sterilized at 121°C at 15 lbs. pressure for 15-20 minutes in an autoclave (Shadwick, 1938).

2.2 Isolation of Pathogen

Diseased leaves were first collected from infected plants and thoroughly washed under running tap water. Along with small healthy tissue diseased portion of the leaves were then cut into small bits and surface sterilized with 1% NaOCI solution for 1 minute followed by washing three times with sterile distilled water and Potato Dextrose Agar media. The petri plates were wrapped and incubated at 27±2°C in an incubator. To avoid bacterial contamination streptomycin @100 ppm, was added to the media at lukewarm stage before pouring Potato Dextrose Agar media into petri plates. Hyphal tip method was used for sub-culturing the fungus in media slants/petri-plates. To obtain pure culture of fungus, single spore technique was used.

2.3 Identification of Pathogen

After incubation of culture for 4-7days, on the basis of morphological characteristics of the colony, mycelium, conidiophores and conidia and their comparison with the available literature, the organism was identified as Alternaria porri. The pathogen (Alternaria porri) isolated produced mycelium. septate Later it produced conidiophores arising singly or in small groups. The conidiophores were straight or flexuous, sometime geniculate, septate, pale or mid brown in colour and measured up to 120 µm long and 6-10 µm thick, with one or several conidial scars. A mature conidiophore usually produced solitary conidium but occasionally it also produced conidia with very short chains, straight or curved, rostrate, beak generally equal to the length of the body of the conidium, pale brown to mid golden brown in colour. Overall length of conidia ranged from 100-300 μ m, 15-20 μ m thick in the broadest part with 7-12 transverse and zero to several longitudinal septa, beak flexuous, pale, 2-4 μ m thick and tapering (Priya *et al.*, 2016).

2.4 Symptoms

The initial symptoms on host leaves and floral stalks are white flecks which enlarge and produce sunken purple lesions sometimes surrounded by a yellow to pale brown border. The disease manifests itself on leaves and seed stalks. Later, the spots enlarge rapidly into purplish areas, further coalescing to form large dead patches coverina several square centimetres of the leaf or shoot area. Their length may go up to 4-6 inches or even more. The purplish area of the spot is separated from the healthy green tissues by a narrow, light-coloured zone. Within a period of 15-21 days alternating become dark zones liaht and clearly differentiated over the whole purple surface of the leaf and seed stalk. A distinct yellowish discoloration usually extends from the spots to the tips and bases of the leaves. The leaves shrivel, usually from the tip (Nolla, 1927).

The most characteristic symptom of this disease as observed is the appearance of dark purple colour on the spots, the dark shade being due to the dark brown colour of the fruiting mycelium. The purple colour has often been seen to fade away in the case of very old lesions, the centre of the spots, however, retain a mild purple colour (Pandotra, 1964).

In order to assess the effect of various treatment, seven treatments applied at different concentrations *viz*. T₁- neem oil (5%), T₂- eucalyptus oil (5%), T₃- clove oil (5%), T₄- castor oil (5%), T₅- mustard oil (5%), T₆- mancozeb 75 WP (0.2%) and T₀- control with each treatment replicating three times to manage the disease and also to evaluate the yield of crop.

Observations were recorded under field condition i.e., disease intensity (%) at 30, 60 and 90 days after incidence of purple blotch and yield.

Percent disease index was calculated by using the following formula (Wheeler, 1969).

Percent Disease Index (PDI) = Sum of individual ratings

Total number of ratings × Maximum disease grade x 100

Pandey et al.; J. Adv. Microbiol., vol. 24, no. 12, pp. 172-179, 2024; Article no.JAMB.128149

Table 1. Disease scale

Grade	Description
0	No disease symptoms.
1	A few spots towards tip covering 10% leaf area.
2	Several dark purplish brown patches covering up to 20% leaf area.
3	Several patches with paler outer zone covering up to 40% leaf area.
4	Leaf streaks covering up to 75% leaf area or breaking of the leaves from centre.
5	Complete drying of the leaves or breaking of the leaves from centre.



Fig. 1. Disease grade scale of purple blotch of garlic



Fig. 2. Field preparation and sowing of seeds

3. RESULTS AND DISCUSSION

3.1 Effects of Essential Oils were Assessed on Plant Disease Intensity and Yield of Garlic

Effects of essential oils were assessed on plant disease intensity of garlic. The data presented in Table 2 and depicted in Fig. 3 revealed that at 30 DAS the disease intensity of purple blotch of garlic significantly decreased in the treatment T_{6} -

Mancozeb (5.75%), followed by T₁- Neem oil (5.98%), T₂- Eucalyptus oil (7.83%), T₃- Clove oil (7.94%), T₄- Castor oil (9.25%) and T₅- Mustard oil (9.85%) as compared to non-treated check, T₀- Control (9.98%).

At 60 DAS the disease intensity of purple blotch of garlic significantly decreased in the treatment T₆- Mancozeb (20.49%) followed by T₁- Neem oil (26.31%), T₃- Clove oil (30.93%), T₂- Eucalyptus oil (31.34%), T₄- Castor oil (32.03%) and T₅-

Mustard oil (33.34%) as compared to non-treated check, T_0 - Control (37.37%).

At 90 DAS the disease intensity of purple blotch of garlic significantly decreased in the treatment T₆- Mancozeb (27.85%) followed by T₁- Neem oil (34.76%), T₃- Clove oil (37.66%), T₂- Eucalyptus oil (37.83%), T₄- Castor oil (39.49%) and T₅- Mustard oil (40.25%) as compared to non-treated check, T₀- Control (49.24%).

Among the treatments, disease intensity (%) recorded in treatments T_{1} - Neem oil @5% (22.35) followed by T_{3} - Clove oil @5% (25.51), T_{4} - Castor oil @5% (26.92) and T_{0} - Control (32.19) showed similar findings (21.35, 25.13, 24.76 and 32.00) respectively supported by Singh *et al.* (2021). T_{2} - Eucalyptus oil @5% (25.66) and T_{5} - Mustard oil @5% (27.81) showed similar finding as 25.44 and 26.10 respectively

supported by Rahmatzai *et al.* (2017). T₆-Mancozeb (18.03) showed similar finding (19.00) supported by Chaurasia *et al.* (2007). The most active component of neem oil is azadirachtin, followed by nimbidol, nimbin, sodium nimbinate, nimbidin, salannin and quercetin. Azadirachtin is a terpene limonoid present in seeds that has properties, which are both antifeedant and toxic to pathogens. Through antimicrobial activity, it inhibits microbial growth or potential to break the cell wall of pathogen (Vijaykumar *et al.*, 2022).

Study of data on yield of garlic is presented in Table 3 and illustrated in Fig. 4 reveals that the yield (t/ha.) significantly increased in the treatment T₆- Mancozeb (5.41) followed by T₁-Neem oil (4.05), T₃- Clove oil (3.83), T₂-Eucalyptus oil (3.73), T₄- Castor oil (3.56) and T₅-Mustard oil (3.31) as compared to non-treated check, T₀- Control (2.61).

Table 2. Effect of treatments on disease intensity (%) of purple blotch of garlic at 30, 60 and 90
DAS

Treatments		Disease intensity (%)			
		30 DAS*	60 DAS [*]	90 DAS*	
T ₀	Control	9.98 ^a	37.37ª	49.24ª	
T ₁	Neem oil (5%)	5.98 ^d	26.31 ^d	34.76 ^d	
T ₂	Eucalyptus oil (5%)	7.83 ^c	31.34°	37.83°	
Тз	Clove oil (5%)	7.94°	30.93°	37.66°	
T_4	Castor oil (5%)	9.25 ^b	32.03°	39.49 ^b	
T ₅	Mustard oil (5%)	9.85 ^{ab}	33.34 ^b	40.25 ^b	
T ₆	Mancozeb (0.2%)	5.75 ^d	20.49 ^e	27.85 ^e	
C.D.	(0.05)	0.65	1.18	1.23	
S.Ed. (±)		0.30	0.54	0.57	

*Mean of three replications



Fig. 3. Effect of treatments on disease intensity of garlic at 30, 60 and 90 DAS

Treatments	Treatment details	Yield (t/ha.)*
To	Control	2.61
T ₁	Neem Oil (5%)	4.05
T ₂	Eucalyptus Oil (5%)	3.73
T ₃	Clove Oil (5%)	3.83
T ₄	Castor Oil (5%)	3.56
T 5	Mustard Oil (5%)	3.31
T ₆	Mancozeb 75 WP (0.2%)	5.41
C.D. (0.05)		0.27
S.Ed. (±)		0.12
i	*Mean of three replication	ons

Table 3. Effect of treatments on yield (t/ha.) of garlic





Among the treatments, yield recorded by the treatments T_1 - Neem oil @5% (4.05) and T_0 -Control (2.61) showed similar findings (4.06 and 2.60) respectively supported by Singh *et al.* (2021). T_6 - Mancozeb (5.41) showed similar finding (5.20) supported by Akter *et al.* (2022).

4. CONCLUSION

Based on the results obtained from present study it was concluded that treatment T1- Neem oil @5% was found most effective against purple blotch disease of garlic. Results of the present study were found to be significantly effective under Prayagraj agro-climatic conditions. It may vary with region and climatic conditions, therefore for validation of the results more such trials should be carried out in future for further recommendations.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative Al technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

ACKNOWLEDGEMENT

The authors are highly thankful and extend sincerest gratitude to the HEAD. their Department of Plant Pathology, Naini Agriculture Institute, Sam Higginbottom University of Aariculture. Technology and Sciences. Prayagraj for all the support and providing all the facilities in completion of the research work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Akter, U. H., Begum, F., Islam, M. R., Prinky, J. N. and Khatun, M. R. (2022). Occurrence of purple blotch disease associated with selected garlic varieties and its management through bio-agents, botanicals and fungicides. *Nepal Journal of Biotechnology*. 10(1): 13-24.
- Chaurasia, C. P. P., Prasad, B. J. and Mandal, A. (2007). Management of leaf blight of garlic with fungicides in Central Tarai of Nepal. *Nepal Agriculture Research Journal.* 8: 63-68.
- Cifferi, R. (1930). Phytopathological survey of Santo Domingo (1925-1929). *Journal of the Department of Agriculture*. 14(4): 30.
- Cooke, M. C. and Ellis, J. B. (1879). *New Jersey fungi*. Grevillea. 8: 11-16.
- Dar, A. A., Sharma, S., Mahajan, R., Mushtaq, M., Salathia, A., Ahamad, S. and Sharma, J. P. (2020). Overview of purple blotch disease and understanding its management through chemical, biological and genetic approaches. *Journal of Integrative Agriculture*. 19(12): 3013-3024.
- Dicklow, M. B. (2013). Storage diseases of onion and garlic. UMass Extension-Vegetable Program.
- FAO (Food and Agriculture Organization). (2021). World food and agriculture -Statistical yearbook 2021. Rome: FAO.
- Harris, J. C., Cottrell, S. L., Plummer, S. and Lloyd, D. (2001). Antimicrobial properties of garlic (*Allium sativum*). *Journal of Applied Microbiology and Biotechnology*. 57: 282-286.
- Kar, S. R., Zacharia, S. and Suresh, B. A. (2021).
 Eco-Friendly Management of Alternaria Leaf Spot of Brinjal (Solanum melongena L.). International Journal of Current Microbiology and Applied Sciences, 10(07): 20-29.
- Kareem, M. A., Murthy, K. V. M. K., Nadaf, H. A. and Waseem, M. A. (2012). Effect of temperature, relative humidity and light on lesion length due to *Alternaria porri* in onion. *Asian Journal of Environmental Science*. 7: 47-49.
- Memane, P. G., Tomar, R. S., Kakade, D. K., Kulkarni, G. U. and Covatia, R. S. (2008).

Effect of clove weight and plant growth regulators on growth and yield of garlic (*Allium sativum* L.). *The Asian Journal of Horticulture*. 3(1): 82-86.

- NHB (National Horticulture Board). (2021-2022). Indian Horticulture Database Ministry of Agriculture, Government of India. http://www.nhb.gov.in.
- Nolla, J. A. B. (1927). A new alternaria disease of onions (*Allium cepa* L.). *Journal of Phytopathology*. 7: 115-132.
- Pandotra, V. R. (1964). Purple blotch disease of onion in Punjab. Studies on the life history, viability and infectivity of the causal organism Alternaria porri. Indian Academy of Sciences. 61: 326-330.
- Prajapati, M. K., Warghat, A. N., Mahajan, S. S. and Simon, S. (2020). Efficacy of organic amendments against the purple blotch of garlic caused by *Alternaria porri* (Ellis) Cif. *Journal of Pharmacognosy and Phytochemistry*. 9(2): 2048-2050.
- Priya, R. U., Sataraddi, A. and Darshan, S. (2016). Survey for purple blotch of onion (*Alternaria porri* (Ellis) Cif.) in northern parts of Karnataka. *International Journal of Agriculture, Environment and Biotechnology*. 9(3): 367-373.
- Rahmatzai, N., Zaitoun, A. A., Madkour, M. H., Ahmady, A., Hazim, Z. and Mousa, M. A. A. (2017). *In vitro* and *in vivo* antifungal activity of botanical oils against *Alternaria solani* causing early blight of tomato. *International Journal of Biosciences*. 10(1): 91-99.
- Saini, S. and Raj, K. (2022). Relative efficacy of fungicides in managing onion purple blotch incited by *Alternaria porri*. *Plant Disease Research*. 37(2): 118-124.
- Shadwick, G. W. Jr. (1938). A study of comparative methods and media used in microbiological examination of creamery butter. *Journal of food science*. 3(3): 287-298.
- Simmons, E.G. (2007). Alternaria: An Identification Manual. CBS Biodiversity Series. CBS Fungal Biodiversity Centre, Utrecht, The Netherlands.
- Singh, S., Tiwari, S., Murmu, J. J. and Jennifer, G. M. J. (2021). Eco-friendly management of purple blotch of garlic (*Allium sativum* L.) caused by *Alternaria porri* (Ellis) Cif. *International Journal of Current Microbiology and Applied Sciences*. 10(3): 1832-1839.
- Tripathi, P. C. and Lawande, K. E. (2006). Cold storage of onion and garlic. *Technical*

Pandey et al.; J. Adv. Microbiol., vol. 24, no. 12, pp. 172-179, 2024; Article no.JAMB.128149

Bulletin No. 15 (2-8). National Research Center for Onion and Garlic, Pune.

Vijaykumar, K. N., Kulkarni, S., Patil, P. V., Kambrekar, D. N. and Shashidhar, T. R. (2022). Eco-friendly management of purple blotch of garlic caused by *Alternaria* porri (Ellis) Ciferri. Biological Forum -An International Journal. 14(2): 1406-1412.

Wheeler, B. E. J. (1969). An introduction to plant diseases. *John Wiley and Sons Ltd.*, London.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/128149